

Brief Insight about Possible Therapeutic Effects of Taurine in Kidney Diseases

Gehad Mohammad Elsayed Ali¹, Abdelmonem Awad Hegazy^{1,2}, Walaa Abdel Haliem Rashad Abdel Haliem¹, Emtethal Mamdouh El-Bestawy¹

1 Human Anatomy and Embryology Department, Faculty of Medicine, Zagazig University, Egypt

2 Human Anatomy and Embryology Department, Faculty of Dentistry, Zarqa University, Jordan

Corresponding author: Gehad Mohammad Elsayed Ali

E-mail: GmElhabashy@medicine.zu.edu.eg, Gehadmuhamnad123@gmail.com,
gehadmohamedelsayed2023@gmail.com

Conflict of interest: None declared

Funding: No funding sources

Abstract

Both the plasma and the cytosol contain high concentrations of taurine, a free amino acid. The kidney is essential in regulating taurine levels. The amino acid shows a peculiar distribution along the nephron, as shown by immunohistochemical analysis. In a number of in vitro and in vivo models, taurine has been shown to have antioxidant properties. It protects glomerular mesangial cells and renal tubular epithelial cells from lipid peroxidation when those cells are cultured in high glucose or low oxygen environments. Animal models of renal illness, such as refractory nephrotic syndrome and diabetic nephropathy, respond favourably to dietary taurine supplementation. Taurine's antioxidant activity is the mechanism by which it exerts its positive effects. Subtotal ablative nephrectomy does not ameliorate acute ischemia or nephrotoxic renal insufficiency or chronic renal failure. More research on taurine's role as a renoprotective drug in experimental renal disease is necessary to completely elucidate its breadth and mechanism of action. The potential of this amino acid as an auxiliary treatment for diabetic nephropathy and progressive glomerulonephritis requires more clinical investigation.

Keywords: Taurine, kidney disease

Tob Regul Sci.™ 2023;9(1): 3891-3901

DOI: doi.org/10.18001/TRS.9.1.273

Introduction

Taurine (2-aminoethanesulfonic acid) is a ubiquitous amino acid in the cells of nearly all body organs. It is present as a free molecule and is not incorporated into structural protein (Chesney, 1985; Huxtable, 1992; Sturman, 1993). The cytosolic concentration of taurine is 1-5 mM depending on the particular type of tissue (Chesney, 1985).

The transmembrane gradient between the plasma and cell can exceed 100, a difference that is maintained by an epithelial Na^+/Cl^- coupled transporter (Huxtable, 1992). Taurine is normally an abundant nutrient in the diet and it is readily absorbed from the gastrointestinal tract (Jacobsen et al., 1968). It is also synthesized from the metabolic precursor, cysteine, via the enzyme, cysteine sulfinic acid decarboxylase (Jacobsen et al., 1968). Although the hepatic biosynthetic capacity for taurine is limited in humans, the effects of dietary taurine deficiency are ameliorated by decreased conjugation of bile acids with taurine and a greater proportion of bile conjugation with glycine (Sturman, 1993). Therefore, taurine is considered a conditionally essential amino acid and serum and tissue levels of the amino acid can be compromised in newborn infants and patients receiving parenteral alimentation with taurine-free solutions for prolonged periods (Gaull, 1989; Zelikovic et al., 1990). Under most conditions, the adequacy of total body taurine balance is determined by the difference between dietary intake and urinary excretion of the amino acid (Chesney, 1985).

The kidney plays a crucial role in regulating body stores of taurine. It possesses the electrogenic Na^+/Cl^- coupled co-transporter in the proximal tubule brush border membrane (Zelikovic et al., 1989). This membrane protein carries out secondary active transport of taurine and other β -amino acids into the cytosol, taking advantage of the sodium gradient generated by the Na^+/K^+ ATPase pump that is present in the basolateral membrane. The activity of this carrier, the V_{max} of the transporter, is directly modulated by nutritional availability of taurine. When there is excessive taurine in the diet, reabsorption is suppressed; in contrast, under conditions of taurine deficiency, the activity of the transporter is upregulated (Chesney et al., 1985). Relative deficiency of this transporter accounts for the amino aciduria and hypertaurinuria observed in newborn infants (Chesney et al., 1986).

Taurine serves many functions including bile acid conjugation, modulation of neurotransmission, stabilization of the retinal membrane and osmoregulation (Chesney, 1985). The accumulation of taurine as a compatible organic osmolyte in cerebral and renal cells under conditions of hyperosmolality is accomplished to preserve cell volume (Trachtman et al., 1988, 1990; Nakanishi et al., 1991). Increased cytosolic taurine content minimizes changes in intracellular ionic strength and perturbations in protein structure and function. These cells enhance the activity of the Na^+/Cl^- coupled cotransporter in response to increases in ambient osmolality. This adaptation has been demonstrated in cerebral cells during chronic hypernatremia and hyperglycemia and in kidney cells during antidiuresis. Cultured MDCK (Madin-Darby canine kidney) cells, analogues of the distal tubule, display osmotically-regulated taurine uptake across the basolateral membrane (Jones et al., 1990; Uchida et al., 1991; Jones et al., 1995). These aspects of taurine handling by the kidney, including response to dietary changes and osmolal disturbances, have been well summarized in a recent report (Jones et al., 1993).

Taurine and oxidant injury

There is considerable information gathered from in vitro reaction systems that taurine is an ineffective antioxidant that is unable to inactivate or scavenge common oxidant species such as the hydroxyl or superoxide radical. It is much less potent than hypotaurine and other sulfur-containing molecules at disrupting the initiation or propagation of oxidation processes (Arouma et al., 1988; Tadolini et al., 1995). Despite the disappointing results that have been collected in chemical studies, there is abundant evidence that taurine functions as an antioxidant in a variety of in vivo biological systems. Taurine reacts with HOCl, a potent oxidizing agent generated by activated neutrophils, to form the less toxic product, taurine chloramine (Thomas et al., 1985; Cantin, 1994). This reaction retards damage to ocular surfaces exposed to HOCl (Nakamori et al., 1993). Taurine protects cultured lymphoblastoid cells against iron-ascorbate or retinol-induced oxidant damage and enhances cell viability (Pasantes-Morales et al., 1984; Pasantes-Morales et al., 1985). Taurine supplementation protects the lung from oxidant-induced damage following exposure to ozone or nitrogen dioxide (Banks et al., 1990; Gordon et al., 1986). In the case of ozone, the pulmonary injury is caused by increased expression of inducible nitric oxide synthase in lung macrophages and increased nitric oxide production. Taurine, via the formation of taurine chloramine, inhibits inducible nitric oxide synthase (iNOS) and prevents lung damage (Cantin, 1994; Schuller-Levis et al., 1994). Taurine alone or in combination with niacin reduces pulmonary fibrosis following intratracheal installation of the oxidants, amiodarone (Wang et al., 1992a) or bleomycin (Wang et al., 1992b). Finally, taurine attenuates adriamycin-induced cardiotoxicity in perfused chick hearts by decreasing myocardial lipid peroxidation (Hamaguchi et al., 1989).

Oxidant injury and the kidney Reactive oxygen molecules are involved in progressive organ damage in several experimental models of kidney disease. In ischemic acute renal failure caused by bilateral renal artery clamping, intra-renal infusion of free radical scavengers such as catalase and superoxide dismutase lessens the reperfusion injury following restoration of renal blood flow (Paller et al., 1984). The hydroxyl radical contributes to the nephrotoxicity of the aminoglycoside antibiotic, gentamicin (Walker et al., 1988).

Oxygen free radicals are implicated in the pathogenesis of the glomerulopathies induced by puromycin aminonucleoside and adriamycin. Administration of allopurinol, catalase and the iron chelating agent, desferrioxamine, reduces the severity of acute phase puromycin aminonucleoside-induced injury (Diamond et al., 1986; Thakur et al., 1988). Provision of tungsten, an inhibitor of renal xanthine oxidase/xanthine dehydrogenase, alleviates adriamycin nephropathy (Ginver et al., 1990). Administration of vitamin E reduces proteinuria, maintains tissue levels of sulfhydryl proteins and lessens structural injury in nephrotoxic serum nephritis (Endreffe et al., 1991). In diabetic nephropathy, hyperglycemia directly promotes lipid peroxidation of cell membranes and causes renal cell injury (Baynes, 1991). Glycosylation of proteins initiates a series of autooxidative chemical reactions

resulting in the accumulation of advanced glycosylation end products (AGEs) in the kidney. These compounds contribute to the development of diabetic nephropathy (Baynes, 1991).

Taurine as a renal antioxidant: In vitro studies

Most of the evidence that taurine is an antioxidant in cultured renal cells is derived from work examining the direct effects of hypoxia or nephrotoxic agents. Addition of taurine (0.1-1.0 mM) to renal transplant preservative solutions prolonged the survival and enhanced the viability of LLC-PK1 cell monolayers that were exposed to hypoxic conditions followed by reoxygenation (Wingenfeld et al., 1994). The protective effect was more apparent with the University of Wisconsin compared to the Euro-Collins perfusion solution. In contrast, Heyman et al. (1992) and Baines et al. (1990) failed to demonstrate any protective effect of adding taurine to the perfusion solution on the extent of medullary tubule damage in the isolated perfused rat kidney preparation. However, glycine was able to reduce the degree of morphologic injury to medullary tubules. Similarly, glycine but not taurine attenuated lethal renal cell injury, assayed by LDH and K⁺ release, in response to increases in cytosolic calcium in the two cultured tubular epithelial cell lines, MDCK and LLC-PK1 (Weinberg et al., 1991a,b). Exposure of cultured glomerular mesangial cells to high glucose concentration (28 mM) increases lipid peroxidation assessed by malondialdehyde or conjugated diene content (Trachtman et al., 1993a). An elevated ambient glucose also inhibits mesangial cell proliferation (Trachtman et al., 1994). Addition of taurine (500 μ M) to the ambient medium prevented these changes (Trachtman et al., 1993a, 1994). The protection against high glucose-induced oxidant injury was reproduced by the addition of vitamin E, consistent with a direct antioxidant effect of the amino acid (Trachtman, 1994).

Taurine as a renal antioxidant: In vivo studies

Localization of taurine within the kidney A prerequisite for understanding the potential function of taurine as a renal antioxidant is clarification of the distribution of the amino acid within the different structural components of the nephron. In the 1980's, a technique was developed for making antibodies to amino acids by conjugating them with glutaraldehyde to proteins such as bovine serum albumin or polylysine (Storm-Mathisen et al., 1983). This advance enabled the localization of amino acids to be determined in a variety of tissues. Using highly specific polyclonal antibodies to taurine that were preabsorbed with structurally similar amino acids, several investigators have examined the localization of taurine within the kidney. Trachtman et al. (1993b) incubated normal rat kidney tissue with an antibody raised to a taurine-glutaraldehyde-BSA conjugate and visualized the amino acid by the biotin-avidin-peroxidase-diaminobenzidine method. They found that the staining intensity for taurine was most pronounced in medullary tubules with minimal secondary staining in proximal tubules and the glomeruli. These findings were independently confirmed by two other groups of investigators. Amiry-Moghaddam et al. (1994) used peroxidase-antiperoxidase and immunogold staining procedures to demonstrate localization of taurine primarily in collecting duct cells, proximal convoluted and straight

tubules and thin descending limbs of Henle. Finally, Ma et al. (1994) utilized an immunoperoxidase method to verify that taurine reactivity was most prominent in glomeruli and the collecting tubules. The heterogeneous staining pattern of taurine within renal cells under conditions of stable dietary intake and urinary osmolality argues in favor of a specific role for this amino acid in the preservation of renal structure and function under normal conditions and in disease states.

Acute and chronic renal failure In rats with acute renal failure induced by bilateral nephrectomy, there is a rise in the plasma taurine concentration; in addition, hepatic taurine content is increased while in skeletal muscle, heart, and brain tissue, the amino acid level is unchanged (Michalk et al., 1983). There are no comparable data about the effect of acute renal failure on plasma or tissue levels of taurine.

In rats with chronic renal failure, the plasma taurine concentration and cerebral content of the amino acid are elevated (Michalk et al., 1983). Patients with chronic renal failure manifest a similar increase in plasma taurine concentration. This change has been documented early in the course of this disease, prior to the onset of protein malnutrition (Ceballos et al., 1990). In patients on hemodialysis, the elevated plasma taurine concentration was normalized by the dialysis treatment (Jung et al., 1991). However, under these circumstances, there was evidence of disparate taurine levels in circulating blood cells. Thus, while taurine concentration was elevated in erythrocytes and normal in granulocytes and lymphocytes, it was lower than normal in platelets (Jung et al., 1991). Skeletal muscle taurine content is diminished compared to normal in patients with chronic renal failure (Bergstrom et al., 1989). The observation that the taurine level in muscle remains low up to 3 months post-renal transplantation is additional evidence in support of body taurine depletion in this condition (Perfumo et al., 1994). Despite the abundant evidence that reactive oxygen molecules contribute to acute and chronic renal failure, we have failed to demonstrate any beneficial effect of taurine under these circumstances. Thus, in rats with bilateral renal ischemia for 30-60 minutes, pre-treatment with a 1% taurine solution for 2 weeks did not reduce the peak serum creatinine concentration observed after 24 hours or hasten recovery of renal function (Trachtman, unpublished observation). In addition, this regimen failed to ameliorate the severity of gentamicin nephrotoxicity induced by injection of the aminoglycoside antibiotic (100mg/kg/day) for 7 days (Trachtman, unpublished observation). Finally, this dietary treatment had no discernible impact on mortality or the long term renal outcome in rats subjected to 5/6 nephrectomy and observed for 8- 10 months (Trachtman et al., 1986).

Nephrotic syndrome Puromycin aminonucleoside (PAN) is an epithelial cell toxin and administration of a single dose of this agent to experimental animals causes a lesion that resembles minimal change nephrotic syndrome (Diamond et al., 1986). Venkatesan et al. (1993, 1994) examined rats given PAN, 100mg/kg intraperitoneally, and reported that provision of taurine, in a daily dose of 500mg for 10 days, reduced total proteinuria, albuminuria and urinary excretion of N-acetyl-fl-D-glucosaminidase by nearly 50%. This treatment also ameliorated the hyperlipidemia observed in these animals. The effects of taurine were associated with restoration of renal content of glutathione, total thiols, ascorbic

acid and vitamin E towards the levels observed in normal rats. These same investigators have studied the effect of taurine on the nephropathy induced by another epithelial cell toxin, adriamycin.

Diabetic nephropathy

Taurine depletion has been noted in nerve tissue isolated from rats with streptozocin (STZ)-induced diabetes for 21 days (Stevens et al., 1993). In 39 patients with insulin-dependent diabetes mellitus, plasma and platelet taurine levels were reduced compared to normal subjects; both values were normalized by taurine supplementation (1.5 g/day) for 90 days (Franconi et al., 1995). In rats with STZ-diabetes, Goodman and Shihabi studied the effect of giving taurine as 0.1% drinking water for 8 weeks (Goodman et al., 1990). Although there was no change in the severity of hyperglycemia or glycosuria, the hypertriglyceridemia and hypercholesterolemia seen in the untreated diabetic animals was reduced by taurine treatment. The duration of the study was not sufficient to elucidate the effect of this maneuver on renal function and structure. We recently reported the effects of long-term dietary taurine supplementation (1% drinking water) for 1 year to rats with STZ-diabetes (Trachtman et al., 1995). This treatment did not ameliorate hyperglycemia. However, from 6 months on, there was a 50% reduction in proteinuria and albuminuria. Although the total kidney and estimated single nephron glomerular filtration rate remained elevated in taurine-treated rats, there was a reversal of glomerular hypertrophy and prevention of glomerulosclerosis and tubulointerstitial fibrosis. These changes occurred in association with reduced serum free iron concentration and lowered renal cortical malondialdehyde content. Finally, taurine supplementation reduced the content of pentosidine and total fluorescence, advanced glycosylation end products, in collagen extracted from rat skin. These changes were paralleled by increased immunohistochemical staining intensity for taurine in glomeruli and medullary tubules (Trachtman et al., 1993b). In contrast to the beneficial effects of taurine on the course of experimental diabetic nephropathy, administration of a diet that was moderately enriched in vitamin E exacerbated the renal lesion in the diabetic rats (Trachtman et al., 1995). The effect of taurine on the course of diabetic nephropathy has not been examined in humans.

Challenges for the future

There is much additional work that is needed to gain a better understanding of the role of taurine in the modulation of renal disease. The mechanism of action of this amino acid requires further clarification. The contribution of taurine acting as an organic osmolyte and the impact of possible changes in the cytosolic content of molecules such as myo-inositol should be addressed in the development of diabetic nephropathy. The interaction of taurine with various intracellular signalling systems and peptide growth factors should be addressed because these are likely to contribute to the biological effects of the amino acid. It is conceivable that the beneficial effect of taurine, acting as an antioxidant, is based upon its ability to regulate inducible or endothelial nitric oxide synthase, alter nitric oxide synthesis, or inhibit the accumulation of oxidant NO products. Alternatively, taurine may act in combination with NF- κ B, an oxidant sensitive transcription factor, or directly modulate oxidant-

sensitive transcription of extracellular matrix genes by virtue of its capacity to alter redox sensitive promoter regions. Finally, like vitamin E, taurine may inhibit activation of protein kinase C and reduce synthesis of extracellular matrix proteins (Kunisaki, 1994). This mechanism may be especially important in the pathogenesis of diabetic nephropathy. All of these effects are amenable to direct study using a combination of in vitro and in vivo systems. The optimal taurine dosage needed to achieve a renoprotective effect and the timing of initiation of amino acid supplementation should be investigated in the relevant disease models. It is necessary to ascertain which particular disease states that are characterized by oxidant-mediated injury are amenable to modification by taurine treatment. The features that define responsiveness to taurine need to be determined. For example, it is unclear why taurine is ineffective in aminoglycoside nephrotoxicity or acute ischemic renal failure while it exerts a beneficial effect in puromycin aminonucleoside nephropathy or diabetic kidney disease. Differences between chronic versus acute disease states may be critical in defining the efficacy of taurine treatment. Finally, consideration should be given to the performance of clinical trials to ascertain whether taurine supplementation can attenuate the severity of refractory nephrotic syndrome or delay the progression of incipient diabetic nephropathy.

Conclusion

Much evidence has developed over the past decade suggesting that taurine improves kidney function and structure in a wide range of disease situations. Most of the evidence indicates that taurine is protecting renal cells by functioning as an antioxidant. More research is needed to understand whether or not taurine has therapeutic potential (diabetic nephropathy vs. 5/6 nephrectomy). Furthermore, it is evident that taurine does not restore renal function. The evidence suggests instead that it acts in a generic fashion to slow the decline of kidney function. Its mode of action is unique, meaning it may work in tandem with other interventions like dietary protein restriction and angiotensin converting enzyme inhibitors to slow the progression of kidney impairment. Existing evidence suggests that taurine may be useful as an additional treatment to reduce kidney injury in a wide range of renal disorders, but more research is needed.

No Conflict of interest.

References:

- [1] Amiry-Moghaddam M, Nagelhus E, Ottersen OP (1994) Light- and electron microscopic distribution of taurine, an organic osmolyte, in rat renal tubule cells. *Kidney Int* 45: 10-22
- [2] Arouma OI, Halliwell B, Hoey BM, Butler J (1988) The antioxidant action of taurine, hypotaurine, and their metabolic precursors. *Biochem J* 256: 251-255
- [3] Baines AD, Shaikh N, Ho P (1990) Mechanisms of perfused kidney cytoprotection by alanine and glycine. *Am J Physiol* 259: F80—F87
- [4] Banks MA, Porter DW, Martin WG, Castranova V (1990) Effects of in vitro ozone exposure on peroxidative damage: membrane leakage and taurine content of rat alveolar macrophages. *Toxicol Appl Pharmacol* 105: 55—

- [5] Baynes JW (1991) Role of oxidative stress in development of complications in diabetes.
- [6] Diabetes 40: 405-412
- [7] Bergstrom J, Alvestrand A, Furst P, Lindholm B (1989) Sulphur amino acids in plasma and muscle in patients with chronic renal failure: evidence for taurine depletion. J Intern Med 226: 189—194
- [8] Brenner BM, Anderson S (1992) The interrelationships among filtration surface area, blood pressure, and chronic renal disease. J Cardiovasc Pharmacol 19 [Suppl] 6: S1— S7
- [9] Cantin AM (1994) Taurine modulation of hypochlorous acid-induced lung epithelial cell injury in vitro: role of anion transport. J Clin Invest 93: 606—614
- [10] Ceballos I, Chauveau P, Guerin V, Bardet J, Parvy P, Kamoun P, Jungers P (1990) Early alterations of plasma free amino acids in chronic renal failure. Clin Chim Acta 188:101—108
- [11] Chesney RW (1985) Taurine: its biological role and clinical implications. Adv Pediatr 32: 1—42
- [12] Chesney RW, Gusowski N, Dabbagh S (1985) Renal cortex taurine content regulates renal adaptive response to altered dietary intake of sulfur amino acids. J Clin Invest 76: 2213—2221
- [13] Chesney RW, Gusowski N, Zelikovic I, Padilla M (1986) Developmental aspects of renal Q-amino acid transport. V. Brush border membrane transport in nursing animals: effect of age and diet. Pediatr Res 20: 890-894
- [14] Culpepper RW, Schoolwerth AC (1992) Remnant kidney oxygen consumption: hypermetabolism or hyperbole? J Am Soc Nephrol 3: 151—156
- [15] Diamond JR, Bonventre JV, Karnovsky MJ (1986) A role for oxygen free radicals in aminonucleoside nephrosis. Kidney Int 29: 478—483
- [16] Endreffe E, Turi S, Laszik Z, Bereczki C, Kasa K (1991) Effects of vitamin E on tissueoxidation in nephrotoxic (anti-glomerular basement membrane) nephritis. Pediatr Nephrol 5: 312—317
- [17] Franconi F, Bennardini F, Mattana A, Miceli M, Ciuti M, Mian M, Gironi A, Anichini R, Seghieri G (1995) Plasma and platelet taurine are reduced in subjects with insulin- dependent diabetes mellitus: effects of taurine supplementation. Am J Clin Nutr 61:1115-1119
- [18] Fujita T, Sato Y (1988) Hypotensive effect of taurine: possible involvement of the sympathetic nervous system and endogenous opiates. J Clin Invest 82: 993— 997
- [19] Fujita T, Sato Y, Ando K (1986) Changes in cardiac and hypothalamic noradrenergic activity with taurine in DOCA-salt rats. Am J Physiol 251: H926—H933
- [20] Fujita T, Ando K, Noda H, Ito Y, Sato Y (1987) Effects of increased adrenomedullary activity and taurine in young patients with borderline hypertension. Circulation 75: 525—532
- [21] Gaull GE (1989) Taurine in pediatric nutrition: review and update. Pediatrics 83: 433—442
- [22] Ginver F, Gusmano R, Oleggini R, et al. (1990) Renal purine efflux and xanthine oxidase activity during experimental nephrosis in rats: difference between puromycin aminonucleoside and adriamycin nephrosis. Clin Sci 78: 283—293
- [23] Goodman HO, Shihabi ZK (1990) Supplemental taurine in diabetic rats: effects on plasma glucose and triglycerides. Biochem Med Metab Biol 43: 1-9
- [24] Gordon RE, Shaked AA, Solano DF (1986) Taurine protects hamster bronchioles from acute N -induced alterations. Am J Pathol 125: 585-600
- [25] Grond J, Weening JJ, Elema JD (1984) Glomerular sclerosis in nephrotic rats: comparison of the long-term effects of adriamycin and aminonucleoside. Lab Invest 51: 277—

- [26] 284
- [27] Hamaguchi T, Azuma J, Harada H, Takahashi K, Kishimoto S, Schaffer SW (1989) Protective effect of taurine against doxorubicin-induced cardiotoxicity in perfused chick hearts. *Pharmacol Res* 21: 729—734
- [28] Heyman S, Spokes K, Rosen S, Epstein FH (1992) Mechanism of glycine protection in
- [29] hypoxic injury: analogies with glycine receptor. *Kidney Int* 42: 41-45 Huxtable RJ (1992) Physiological actions of taurine. *Physiol Rev* 72: 101—163
- [30] Ideishi M, Miura S, Sakai T, Sasaguri M, Misumi Y, Arakawa K (1994) Taurine amplifies renal kallikrein and prevents salt-induced hypertension in Dahl rats. *J Hypertens* 12:653—661
- [31] Jacobsen JG, Smith LH Jr (1968) Biochemistry and physiology of taurine and taurine
- [32] derivatives. *Physiol Rev* 48: 424—511
- [33] Johnson RJ (1994) The glomerular response to injury: progression or resolution? *Kidney Int* 45: 1769—1782
- [34] Jones DP, Miller LA, Chesney RW (1990) Adaptive regulation of taurine transport in two
- [35] continuous renal epithelial cell lines. *Kidney Int* 38: 219—226
- [36] Jones DP, Chesney RW (1993) Polarized nature of taurine transport in LLC-PK, and MDCK cells: further characterization of divergent transport models. *Amino Acids* 5: 329—339
- [37] Jones DP, Miller LA, Chesney RW (1995) The adaptive roles of external taurine concentration and medium osmolality in the regulation of taurine transport in LLC-PK, and MDCK cells. *Pediatr Res* 37: 227—232
- [38] Jung BC, Laidlaw SA, Kopple JD (1991) Taurine levels in plasma and blood cells in patients undergoing routine maintenance hemodialysis. *Am J Kidney Dis* 18: 74— 79
- [39] Klahr S, Schreiner G, Ichikawa I (1988) The progression of renal disease. *N Engl J Med*
- [40] 318: 1657—1666
- [41] Kunisaki M, Bursell SE, Umeda F, Nawata H, King GL (1994) Normalization of diacylglycerol-protein kinase C activation by vitamin E in aorta of diabetic rats and cultured rat smooth muscle cells exposed to elevated glucose levels. *Diabetes* 43: 1372—1377
- [42] Li JZ, Sharma R, Dileepan KN, Savin VJ (1994) Polymorphonuclear leukocytes increase glomerular albumin permeability via hypohalous acid. *Kidney Int* 46: 1025— 1030
- [43] dia N, Aoki E, Semba R (1994) An immunohistochemical study of aspartate, glutamate and taurine in rat kidney. *J Histochem Cytochem* 42: 621—626
- [44] Michalk DV, Essich HJ, Bohles HJ, Scharer K (1983) Taurine metabolism in experimental renal failure. *Kidney Int* 24 [Suppl]: 516—S21
- [45] Nakamori K, Koyama I, Nakamura T, Nemoto M, Yoshida T, Umeda M, Inoue K (1993) Quantitative evaluation of the effectiveness of taurine in protecting the ocular surface against oxidant. *Chem Pharm Bull* 41: 335—338
- [46] Nakanishi T, Uyama O, Sugita M (1991) Osmotically regulated taurine content in rat renal inner medulla. *Am J Physiol* 261: F957—F962
- [47] Nankivell BJ, Tay YC, Boadle RA, Harris DCH (1994) Dietary protein alters tubular iron accumulation after partial nephrectomy. *Kidney Int* 45: 1006—1013
- [48] Nara Y, Yamori Y, Lovenberg W (1978) Effect of dietary taurine on blood pressure in spontaneously hypertensive rats. *Biochem Pharmacol* 27: 2689-2692
- [49] Paller MS, Hoidal JR, Ferris TF (1984) Oxygen free radicals in ischemic acute renal failure in the rat. *J Clin Invest* 74: 1156-1164

- [50] Pasantes-Morales H, Wright CE, Gaull GE (1984) Protective effect of taurine, zinc, and tocopherol on retinol-induced damage in human lymphoblastoid cells. *J Nutr* 114: 2256—2261
- [51] Pasantes-Morales H, Wright CE, Gaull GE (1985) Taurine protection of lymphoblastoid cells from iron-ascorbate induced damage. *Biochem Pharmacol* 34: 2205—2207
- [52] Perfumo F, Canepa A, Divino-Filho JC, Nilsson E, Carrea A, Verrina E, Gusmano R, Bergstrom J (1994) Muscle and plasma amino acids and nutritional status in kidney- transplanted children. *Nephrol Dial Transplant* 9: 1778—1785
- [53] Schuller-Levis G, Quinn MR, Wright C, Park E (1994) Taurine protects against oxidant- induced lung injury: possible mechanism(s) of action. In: Huxtable R (ed) *Taurine in health and disease*. Plenum Press, New York, pp 31-39
- [54] Stevens MJ, Lattimer SA, Kamijo M, Van Huysen C, Sima AAF, Greene DA (1993) Osmotically-induced nerve taurine depletion and the compatible osmolyte hypothesis in experimental diabetic neuropathy in the rat. *Diabetologia* 36: 608—614
- [55] Storm-Mathisen J, Leknes AK, Bore AT, Vaaland JL, Edminson P, Haug FS, Ottersen OP (1983) First visualization of glutamate and GABA in neurones by immuno- cytochemistry. *Nature*: 517-520
- [56] Sturman JA (1993) Taurine in development. *Physiol Rev* 73: 119—147
- [57] Tadolini B, Pintus G, Pinna GG, Bennardini F, Franconi F (1995) Effects of taurine and hypotaurine on lipid peroxidation. *Biochem Biophys Res Commun* 213: 820— 826
- [58] Thakur V, Walker PD, Shah SV (1988) Evidence suggesting a role for hydroxyl radical in puromycin aminonucleoside-induced proteinuria. *Kidney Int* 34: 494-499
- [59] Thomas EL, Grisham MB, Melton DF, Jefferson MM (1985) Evidence for a role of taurine in the in vitro oxidative toxicity of neutrophils toward erythrocytes. *J Biol Chem* 260: 3321—3329
- [60] Trachtman H (1994) Vitamin E prevents glucose-induced lipid peroxidation and in- creased collagen production in cultured rat mesangial cells. *Microvasc Res* 47: 232— 239
- [61] Trachtman H, Del Pizzo R (1989) Taurine does not afford protection against progressive renal failure in the 5/6 nephrectomy model of chronic renal disease. *Am J Kidney Dis* 14: 438 (abstract)
- [62] Trachtman H, Barbour R, Sturman JA, Finberg L (1988) Taurine is a cerebral osmoprotective molecule, “no longer idiogenic”, in chronic hypernatremic dehydra-tion. *Pediatr Res* 23: 35-39
- [63] Trachtman H, Del Pizzo R, Rao PS, Rujikarn N, Sturman JA (1989) Taurine lowers blood pressure in the spontaneously hypertensive rat by a catecholamine independent mechanism. *Am J Hypertens* 2: 909—912
- [64] Trachtman H, Del Pizzo R, Sturman JA (1990) Taurine and osmoregulation III. Taurine deficiency protects against cerebral edema during acute hyponatremia. *Pediatr Res* 27: 85—88
- [65] Trachtman H, Wilson D, Rao PS (1992a) The role of oxygen free radicals in the develop-
- [66] ment of chronic renal failure. *Life Sci* 50: 1877-1883
- [67] Trachtman H, Del Pizzo R, Futterweit S, Levine D, Rao PS, Valderrama E, Sturman JA(1992b) Taurine attenuates renal disease in chronic puromycin aminonucleoside nephropathy. *Am J Physiol* 262: F117—F123
- [68] Trachtman H, Futterweit S, Bienkowski RS (1993a) Taurine prevents glucose-induced lipid peroxidation and increased collagen production in cultured rat mesangial cells. *Biochem Biophys Res Commun* 191: 759-765
- [69] Trachtman H, Lu P, Sturman JA (1993b) Immunohistochemical localization of taurine in rat renal tissue: studies in experimental disease states. *J Histochem Cytochem* 41: 1209—1216
- [70] Trachtman H, Futterweit S, Prenner J, Hanon S (1994) Antioxidants reverse the antiproliferative effect of high glucose and advanced glycosylation end products in cultured rat mesangial cells. *Biochem Biophys Res*

- [71] Trachtman H, Futterweit S, Maesaka J, Ma C, Valderrama E, Fuchs A, Tarectecan AA, Rao PS, Sturman JA, Boles TH, Fu MX, Baynes J (1995) Taurine ameliorates chronic streptozocin-induced diabetic nephropathy in rats. *Am J Physiol* 269: F429—F438
- [72] Ucida S, Nakanishi T, Kwon HM, Preston AS, Handler JS (1991) Taurine behaves as an osmolyte in Madin-Darby canine cells: protection by polarized, regulated transport of taurine. *J Clin Invest* 88: 656-662
- [73] Venkatesan N, Rao PV, Arumugam V (1993) Inhibitory effect of taurine on puromycinaminonucleoside-induced hyperlipidemia in rats. *J Clin Biochem Nutr* 15: 203—210
- [74] Venkatesan N, Rao PV, Arumugam V (1994) In vivo protective effect of taurine on puromycin aminonucleoside-induced nephrosis in rats. *J Clin Biochem Nutr* 16: 91— 100
- [75] Venkatesan N, Karthikeyan J, Arumugam V: Protection by taurine against adriamycin-induced hyperlipidemia and proteinuria in rats. *Proc Soc Exp Biol Med* (under re- view)
- [76] Walker PD, SV Shah (1988) Evidence suggesting a role for hydroxyl radical in gentamicin-induced acute renal failure. *J Clin Invest* 81: 334-341
- [77] Wang O, Hollinger MA, Giri SN (1992a) Attenuation of amiodarone-induced lung fibrosis and phospholipidosis in hamsters by taurine and/or niacin treatment. *J Pharmacol Exp Therap* 262: 127-132
- [78] Wang Q, Hyde DM, Giri SN (1992b) Abatement of bleomycin-induced increases in vascular permeability, inflammatory cell infiltration, and fibrotic lesions in hamster lungs by combined treatment with taurine and niacin. *Lab Invest* 67: 234—241
- [79] Weinberg JM, Venkatachalam MA, Roeser h'F, Davis JA, Varani J, Johnson KJ (1991a) Amino acid protection of cultured kidney tubule cells against calcium ionophore- induced lethal cell injury. *Lab Invest* 65: 671-678
- [80] Weinberg JM, Nissim I, Roeser NF, Davis JA, Schultz S, Nissim I (1991b) Relationships between intracellular amino acid levels and protection against injury to isolated proximal tubules. *Am J Physiol* 260: F410—F419
- [81] Wingenfeld P, Strubind S, Gehrmann U, Minor T, Isselhard W, Michalk D (1994) Protecting effect of taurine against hypoxic cell damage in renal tubular cells cultured in different transplant preservation solutions. *Adv Exp Med Biol* 359: 159—169
- [82] Zelikovic I, Stejskal-Lorenz E, Lohstroh P, Budreau A, Chesney RW (1989) Anion dependence of taurine transport by rat renal brushborder membrane vesicles. *Am J Physiol* 256: F646—F655
- [83] Zelikovic I, Chesney RW, Friedman AL, Ahlfors CE (1990) Taurine depletion in very low birth weight infants receiving prolonged total parenteral nutrition: role of renal immaturity. *J Pediatr* 116: 301-306