

# Protective Effects of Butylphthalide on Cerebral Infarction Induced by Local Ischemic Injury and Regulation Mechanism of the Pi3k/Akt/Gsk-3 $\beta$ Signaling Pathway

Chong Liu  
Yan Cao

Chong Liu, Department of Neurology, Nanyang First People's Hospital, Nanyang 473000, Henan Province, China, Yan Cao, Department of Neurology, Taikang Tongji (Wuhan) Hospital, Wuhan 430050, Hubei Province, China, \*Corresponding author: Yan Cao, Email: (807385155@qq.com)

**Aim:** We aimed to study the protective effects of butylphthalide (NBP) on cerebral infarction induced by local ischemic injury and regulation mechanism of the PI3K/Akt/GSK-3 $\beta$  signaling pathway. **Materials and methods:** One hundred male Wistar rats aged 12-15 weeks were randomly divided into 5 groups (n=20). The middle cerebral artery occlusion (MCAO) model was established. NBP, P13K specific inhibitor LY294002 and NBP plus LY294002 groups were intraperitoneally administered on the first day after modeling, once a day for 7 days. Sham operation (Sham) and model groups were intraperitoneally given equal amounts of normal saline. Modified neurological severity (mNS) was scored 30 min after administration on the 7th day, and cerebral infarction volume was measured by magnetic resonance imaging. Neuronal damage was detected by Nissl staining. Intact neurons were counted under light microscope. The protein expressions of Akt, P-Akt, GSK-3 $\beta$  and P-GSK-3 $\beta$  were detected by Western blotting. **Results:** The mNS score of NBP group decreased significantly compared with that of model group (P<0.05). Compared with model group, the cerebral infarction volume of NBP group significantly reduced (P<0.05). Compared with model group, the number of intact neurons in NBP group significantly increased (P<0.05). Compared with model group, the phosphorylation levels of Akt and GSK-3 $\beta$  in NBP group significantly increased (P<0.05). **Conclusions:** By activating the PI3K/Akt/GSK-3 $\beta$  signaling pathway, NBP relieves neurological function damage and protects against cerebral infarction induced by local ischemic injury.

**Keywords:** Butylphthalide; PI3K; Akt; GSK-3 $\beta$ ; apoptosis; ischemic injury  
*Tob Regul Sci.*™ 2021;7(5): 1578-1584  
DOI: [doi.org/10.18001/TRS.7.5.80](https://doi.org/10.18001/TRS.7.5.80)

Ischemic cerebral infarction is a clinically common central nervous system disease with high mortality and disability rates, which seriously threatens human health and quality of life<sup>1</sup>. With a complicated pathophysiological process, this disease is a blood flow disorder in the brain that is caused by various factors, leading to ischemia and hypoxia, apoptosis of neurons, functional damage of tissues and cells, as well as neurological dysfunction such as movement, language, sensation and memory<sup>2</sup>. During cell apoptosis, the PI3K/Akt signaling pathway plays a key role in the regulation of related proteins, allowing membrane receptor signal to be

transduced to cells for maintaining cell proliferation and inhibiting apoptosis<sup>3</sup>. As a substrate for Akt,

GSK-3 $\beta$  is a key component regulating cell apoptosis<sup>4</sup>. Butylphthalide (NBP) has significant therapeutic effects on the neurological function of patients with ischemic cerebral infarction. It can enhance the blood circulation in the brain and prevent cerebral infarction caused by ischemia. Meanwhile, NBP improves the energy metabolism in ischemic area and participates in the inhibition of neuronal apoptosis<sup>5</sup>. The mechanism by which NBP protects against cerebral infarction induced by

local ischemic injury by regulating the PI3K/Akt/GSK-3 $\beta$  signaling pathway remains unclear. Therefore, we herein established a rat model of middle cerebral artery occlusion (MCAO) to assess the protective effects of NBP on cerebral infarction induced by local ischemic injury, and the regulation of the PI3K/Akt/GSK-3 $\beta$  signaling pathway. The findings provide valuable experimental evidence for clinical practice.

## MATERIALS AND METHODS

### Experimental animals, reagents and apparatus

SPF-grade healthy Wistar male rats aged 12-15 weeks and weighing 200-220 g were purchased from Beijing Vital River Laboratory Animal Technology Co., Ltd. (China). NBP was bought from Shijiazhuang Pharmaceutical Group NBP Pharmaceutical Co., Ltd. (China). LY294002 was obtained from Sigma (USA). BCA protein concentration detection kit, DAB chemiluminescence kit, and antibodies against Akt, P-Akt, GSK-3 $\beta$  and P-GSK-3 $\beta$  were provided by Shanghai Renjie Biotechnology Co., Ltd. (China). Magnetic resonance imaging (MRI) scanner was purchased from Siemens (Germany). Gel imaging system was bought from Bio-Rad (USA). BXM-950 optical microscope was obtained from Shanghai Bingyu Optical Instrument Co., Ltd. (China).

### Model establishment and grouping

One hundred rats were numbered and then randomly divided into 5 groups (n=20): sham operation (Sham) group, model group, NBP (10 mg/kg) group, P13K specific inhibitor LY294002 (LY, 10 mg/kg) group and NBP (10 mg/kg) plus LY294002 (10 mg/kg) (NBP + LY) group. Model establishment: Under anesthesia, a nylon thread with 0.285 mm diameter was used to occlude blood flow in the anterior segment of MCA and its lateral branch for 2 h, causing local MCA ischemia without affecting blood flow in the anterior cerebral artery<sup>6</sup>. After the nylon thread was withdrawn and the MCA blood flow was restored, the rat MCAO model was successfully established. For the Sham group, only vascular ligation or occlusion was not performed. Each group was

intraperitoneally injected with corresponding drugs on the first day after modeling, once a day for 7 days. The same amounts of normal saline were intraperitoneally given to the Sham and model groups.

### Scoring of modified neurological severity (mNS)

Thirty minutes after administration on the 7th day, the rat behaviors were evaluated by mNS scoring according to the criteria below. 0 point: Rat crawls normally without asymmetric movement; 1 point: forelimb or hind limbs are bent when the tail is lifted vertically; 2 points: rat fails to walk straight based on the criterion for 1 point; 3 points: rat circles leftward while crawling; 4 points: rat falls down leftward during free movement; 5 points: rat drags its left forepaw backward based on the criterion for of 4 points; 6 points: rat fails to support its body or to crawl by itself.

### Measurement of cerebral infarction volume by MRI

After mNS scoring, 6 rats were randomly selected. The cerebral infarction volume was detected by MRI scanner. The rats were placed in the supine position. After the standard axial position was found, the coronal plane was subjected to three-layer scan based on cross-sectional T2-weighted imaging, with a layer thickness of 1.5 mm and a spacing of 0.2 mm. T2-weighted imaging was carried out to measure the volume of cerebral infarction. The infarct area was pale white, and the normal brain tissue area was gray. Cerebral infarction volume (%) = (infarct volume/volume of whole brain tissue)  $\times$  100%. The calculation was conducted using ImageJ software.

### Detection of neurons in brain tissue by Nissl staining

After mNS scoring, 6 rats were randomly selected. Under anesthesia, the chest and abdomen were incised, and the exposed left ventricle was intubated. 4% Paraformaldehyde was perfused until the heart turned white. Then the brain tissue was taken out, placed on ice, immediately fixed with 4% paraformaldehyde, dehydrated under vacuum conditions with gradient concentrations of

ethanol solutions, transparentized with xylene and paraffin-embedded. The treated brain tissue was thereafter sliced into 5  $\mu$ m-thick coronal sections that were subjected to Nissl staining. Six different visual fields were observed under an optical microscope at the magnification of 400 $\times$ , and images were processed by Image-Pro 6.2 software to count intact neurons in the cerebral cortex on the ischemic side.

#### Detection of Akt, P-Akt, GSK-3 $\beta$ and P-GSK-3 $\beta$ protein expressions by Western blotting

After mNS scoring, 6 rats were randomly selected. Under anesthesia, they were killed by cervical dislocation, from which the skull was rapidly disconnected to collect brain tissue on the ischemic side. Subsequently, the cerebral cortex was separated in an ice bath and stored in liquid nitrogen. The brain tissue was homogenized with RIPA lysis buffer, left still in ice bath for 5 min and centrifuged at 13000 rpm and 4°C for 10 min to collect the supernatant. Protein concentration in the supernatant was measured by BCA protein quantification kit. Afterwards, protein samples were resolved by 10% SDS-PAGE, and the product was electronically transferred onto a PVDF membrane. Then the membrane was blocked by 5% TBST at room temperature, incubated overnight with primary antibodies (1:1000 diluted) at 4°C, washed by TBST for 10 min, incubated with HRP-labeled goat anti-rabbit IgG secondary antibody (1:10000 diluted) at room temperature for 2 h, developed by DAB solution and observed by gel imaging analyzer. The relative expression levels of target proteins were detected by using GAPDH as the internal reference.

#### Statistical analysis

All data were analyzed by SPSS16.0 software. The categorical data conforming to normal distribution were expressed as mean  $\pm$  standard deviation. Multigroup comparisons were performed by one-way analysis of variance, and pairwise comparisons were conducted with the independent t test.  $P < 0.05$  was considered statistically significant.

## RESULTS

### Effects of NBP on neurological function of MCAO rats

Compared with the Sham group, the mNS score of the model group was significantly higher ( $P < 0.05$ ). After NBP treatment, the mNS score of the NBP group decreased significantly compared with that of the model group ( $P < 0.05$ ), but the scores of LY and NBP + LY groups were similar to that of the model group ( $P > 0.05$ ) (Fig. 1), indicating that LY294002 blocked NBP from improving the neurological function of MCAO rats.

### Effects of NBP on cerebral infarction volume of MCAO rats

No infarct area was found in the brain tissue of the Sham group. Compared with the Sham group, the cerebral infarction volume of the model group significantly increased ( $P < 0.05$ ). Compared with the model group, the cerebral infarction volume of the NBP group significantly reduced ( $P < 0.05$ ), whereas the volumes of LY and NBP + LY groups were not significantly different ( $P > 0.05$ ) (Fig. 2), suggesting that LY294002 inhibited NBP from repairing ischemic cerebral injury.

### Effects of NBP on cerebral neuronal integrity of MCAO rats

The Sham group had normal neuron morphology. Compared with the Sham group, the model group had abnormal structures of neurons and significantly decreased number of intact neurons ( $P < 0.05$ ). Compared with the model group, the number of intact neurons in the brain tissue of the NBP group significantly increased ( $P < 0.05$ ), but the numbers of LY and NBP + LY groups were not significantly different ( $P > 0.05$ ) (Fig. 3), indicating that LY294002 suppressed NBP from protecting the neurons undergoing ischemic brain injury.

### Effects of NBP on expressions of Akt, P-Akt, GSK-3 $\beta$ and P-GSK-3 $\beta$ in brain tissues of MCAO rats

Western blotting showed that compared with

the Sham group, the phosphorylation levels of Akt and GSK-3 $\beta$  in the model group significantly decreased ( $P < 0.05$ ). Compared with the model group, such levels in the brain tissue of the NBP group significantly increased ( $P < 0.05$ ), whereas the levels of LY and NBP + LY groups were not significantly different ( $P > 0.05$ ) (Fig. 4), revealing that LY294002 counteracted the therapeutic effects of NBP by decreasing the levels of in brain tissues with ischemic injury.

## DISCUSSION

The brain is one of the most complex organs in human body, strongly depending on blood supply. When the arteries supplying blood to the brain are narrowed or blocked, brain tissues undergo ischemia or even necrosis due to insufficient blood and oxygen supplies. Brain function is impaired, which has serious adverse effects on physiological activities<sup>7</sup>. Ischemic cerebral infarction, cardiovascular disease and malignancy are all fatal diseases that pose a great threat to human life<sup>8</sup>.

NBP has been widely used as a therapeutic drug for ischemic cerebrovascular disease in clinical practice, with remarkable effects. In this study, a MCAO model was first established, and then the concentration of butylphthalide was selected based on preliminary experiments. Compared with the model group, NBP-treated rats had significantly reduced volume of cerebral infarction, alleviated neurological damage and decreased number of apoptotic neurons, verifying that cerebral ischemic injury was alleviated. The results are consistent with a previous literature<sup>9</sup>, confirming the therapeutic effects of NBP again.

The PI3K/Akt signaling pathway plays crucial roles in cell migration, mobilization, differentiation and apoptosis resistance<sup>10</sup>. As a key target protein, PI3K promotes the phosphorylation of GSK-3 $\beta$  in downstream pathways by phosphorylating Akt through PDK1<sup>11</sup>. Subsequently, activated GSK-3 $\beta$  inhibits cellular oxidative stress and inflammatory response, and also predominantly resists apoptosis<sup>12,17</sup>.

By enhancing the activity of mitochondrial ATPase, NBP protects mitochondria from ischemic damage, thereby suppressing cell apoptosis<sup>13</sup>. In

addition, NBP may increase the expressions of VEGF and bFGF by acting on the ischemic site of the brain to protect against damage<sup>14</sup>. However, the mechanism remains largely unknown. As a PI3K inhibitor, LY294002 specifically inhibits the activity of PI3K110 subunit and blocks PI3K-mediated signaling pathway<sup>15</sup>. In this study, LY294002 was used to clarify the mechanism by which NBP regulated the PI3K/Akt/GSK3 $\beta$  signaling pathway to protect against local brain injury. The phosphorylation levels of Akt and GSK-3 $\beta$  were significantly augmented in the NBP group compared with those of the model group, and the levels of the LY+ NBP group decreased significantly compared with those of the NBP group. Accordingly, LY294002 reversed NBP-mediated phosphorylation of Akt and GSK3 $\beta$ , indicating that NBP allowed Akt and GSK3 $\beta$  phosphorylation through PI3K. It has previously been reported that the PI3K/Akt/GSK-3 $\beta$  signaling pathway inhibited cell apoptosis after reperfusion injury in the brain, and participated in the repair process thereafter, playing a protective role<sup>16</sup>. Compared with the NBP group, the cerebral infarction volume of the LY + NBP group was significantly enlarged, neurological damage was aggravated, and the number of apoptotic neurons was increased, suggesting that NBP may protect against cerebral ischemic injury by regulating the PI3K/Akt/GSK-3 $\beta$  signaling pathway.

## CONCLUSION

In summary, by activating the PI3K/Akt/GSK-3 $\beta$  signaling pathway, NBP relieves neurological function damage and protects against cerebral infarction induced by local ischemic injury. Nevertheless, whether other pathways are involved still needs further in-depth studies.

## CONFLICT OF INTEREST

The authors declare they have no potential conflicts of interest concerning drugs, products, or services used in the study.

## REFERENCES

1. Yu Z-H, Cai M, Xiang J, et al. PI3K/Akt pathway contributes to neuroprotective effect of Tongxinluo against focal cerebral ischemia and reperfusion injury in rats. *Journal of Ethnopharmacology*. 2016/04/02/2016;181:8-19. doi:<https://doi.org/10.1016/j.jep.2016.01.028>
2. Zhang Q, An R, Tian X, et al.  $\beta$ -Caryophyllene Pretreatment Alleviates Focal Cerebral Ischemia-Reperfusion Injury by Activating PI3K/Akt Signaling Pathway. *Neurochemical Research*. 2017/05/01 2017;42(5):1459-1469. doi:<https://doi.org/10.1007/s11064-017-2202-3>
3. Yao R-Q, Qi D-S, Yu H-L, Liu J, Yang L-H, Wu X-X. Quercetin Attenuates Cell Apoptosis in Focal Cerebral Ischemia Rat Brain Via Activation of BDNF-TrkB-PI3K/Akt Signaling Pathway. *Neurochemical Research*. 2012/12/01 2012;37(12):2777-2786. doi:<https://doi.org/10.1007/s11064-012-0871-5>
4. Wei Y, Hong H, Zhang X, et al. Salidroside Inhibits Inflammation Through PI3K/Akt/HIF Signaling After Focal Cerebral Ischemia in Rats. *Inflammation*. 2017/08/01 2017;40(4):1297-1309. doi:<https://doi.org/10.1007/s10753-017-0573-x>
5. Xu J, Huai Y, Meng N, et al. l-3-n-Butylphthalide Activates Akt/mTOR Signaling, Inhibits Neuronal Apoptosis and Autophagy and Improves Cognitive Impairment in Mice with Repeated Cerebral Ischemia-Reperfusion Injury. *Neurochemical Research*. 2017/10/01 2017;42(10):2968-2981. doi:<https://doi.org/10.1007/s11064-017-2328-3>
6. Chi OZ, Mellender SJ, Kiss GK, Liu X, Weiss HR. Blood-brain barrier disruption was less under isoflurane than pentobarbital anesthesia via a PI3K/Akt pathway in early cerebral ischemia. *Brain Research Bulletin*. 2017/05/01/2017;131:1-6. doi:<https://doi.org/10.1016/j.brainresbull.2017.02.007>
7. Qi Y DD, Jiang H et al. Arctigenin attenuates learning and memory deficits through PI3K/Akt/GSK-3 $\beta$  pathway reducing tau hyperphosphorylation in A $\beta$ -induced AD mice. *Planta Med* 2017;83(1-02)doi:<https://doi.org/10.1055/s-0042-107471>
8. Sun Y, Cheng X, Wang H, et al. dl-3-n-butylphthalide promotes neuroplasticity and motor recovery in stroke rats. *Behavioural Brain Research*. 2017/06/30/2017;329:67-74. doi:<https://doi.org/10.1016/j.bbr.2017.04.039>
9. Bai Q, Song D, Gu L, Verkhatsky A, Peng L. Bi-phasic regulation of glycogen content in astrocytes via Cav-1/PTEN/PI3K/AKT/GSK-3 $\beta$  pathway by fluoxetine. *Psychopharmacology*. 2017/04/01 2017;234(7):1069-1077. doi:<https://doi.org/10.1007/s00213-017-4547-3>
10. Zhu Z, Sun H, Ma G, et al. Bufalin Induces Lung Cancer Cell Apoptosis via the Inhibition of PI3K/Akt Pathway. *International Journal of Molecular Sciences*. 2012;13(2):2025-2035.
11. Zhang C, Zhao S, Zang Y, et al. The efficacy and safety of Dl-3n-butylphthalide on progressive cerebral infarction: A randomized controlled STROBE study. *Medicine (Baltimore)*. 2017;96(30):e7257-e7257. doi:<https://dx.doi.org/10.1097%2FMD.00000000000007257>
12. Wang H, Li Y, Wu Q, Xu C, Liu Q. Combination of butylphthalide with umbilical mesenchymal stem cells for the treatment of delayed encephalopathy after carbon monoxide poisoning. *Medicine (Baltimore)*. 2016;95(49):e5412-e5412. doi:<https://dx.doi.org/10.1097%2FMD.00000000000005412>
13. Zhu J, Wu D, Zhao C, et al. Exogenous Adipokine Peptide Resistin Protects Against Focal Cerebral Ischemia/Reperfusion Injury in Mice. *Neurochemical Research*. 2017/10/01 2017;42(10):2949-2957. doi:<https://doi.org/10.1007/s11064-017-2326-5>
14. Wang C-Y, Wang Z-Y, Xie J-W, et al. Dl-3-n-butylphthalide-induced upregulation of antioxidant defense is involved in the enhancement of cross talk between CREB and Nrf2 in an Alzheimer's disease mouse model. *Neurobiology of Aging*. 2016/02/01/ 2016;38:32-46. doi:<https://doi.org/10.1016/j.neurobiolaging.2015.10.024>
15. Avni D, Glucksam Y, Zor T. The Phosphatidylinositol 3-kinase (PI3K) inhibitor LY294002 modulates cytokine expression in macrophages via p50 nuclear factor kappa B inhibition, in a PI3K-independent mechanism. *Biochemical Pharmacology*. 2012/01/01/ 2012;83(1):106-114. doi:<https://doi.org/10.1016/j.bcp.2011.09.025>
16. Xu Z-P, Gan G-S, Liu Y-M, et al. Adiponectin Attenuates Streptozotocin-Induced Tau Hyperphosphorylation and Cognitive Deficits by Rescuing PI3K/Akt/GSK-3 $\beta$  Pathway. *Neurochemical Research*. 2018/02/01 2018;43(2):316-323. doi:<https://doi.org/10.1007/s11064-017-2426-2>
17. Pallavi Sharad Suryarao, Namrata Nishant Gaonkar1 , Prerana Vinod Hariyani, Darshan Wable NDS. Comparison between Oral Health Status of Institutionalised and Home Stay Disabled Children in Western Maharashtra Region. *J Nat Sci Biol Med*. 2020;11(2):179-182. doi:10.4103/jnsbm.JNSBM

Figure. 2.

Effects of NBP on cerebral infarction volume of MCAO rats. \*Compared with Sham group,  $P < 0.05$ ; #compared with model group,  $P < 0.05$ . A: Sham group; B: model group; C: NBP group; D: LY group; E: NBP + LY group.




Figure. 3.

Nissl staining results for effects of NBP on cerebral neuronal integrity of MCAO rats (number of neurons in a  $\times 400$  visual field). \*Compared with Sham group,  $P < 0.05$ ; #compared with model group,  $P < 0.05$ . A: Sham group; B: model group; C: NBP group; D: LY group; E: NBP + LY group.




Figure. 4.

Effects of NBP on expressions of Akt, P-Akt, GSK-3 $\beta$  and P-GSK-3 $\beta$  in brain tissues of MCAO rats. \*Compared with Sham group,  $P < 0.05$ ; #compared with model group,  $P < 0.05$ .

